

Does Smoothing Matter?

Martin Roček

*C.N. Yang Institute for Theoretical Physics
SUNY, Stony Brook, NY 11794-3840, USA*

and

*Institute for Theoretical Physics
University of Amsterdam, Valckenierstraat 65
1018 XE Amsterdam, the Netherlands
rocek@insti.physics.sunysb.edu*

Patrick van Nieuwenhuizen

*Ward Melville High School
180 Old Town Road, East Setauket, NY 11790
PatvanN@gmail.com*

Abstract

We study how inhomogeneities modify large scale parameters in General Relativity. For a particular model, we obtain exact results: we compare an infinite string of extremal black holes to a corresponding smooth line with the same mass and charge in five dimensions. We find that the effective energy density does not differ significantly.

1 Introduction

Less than 5% of all matter and energy in the universe is visible—the origin of the rest is a mystery. Dark Energy (DE) constitutes perhaps 70% of this missing mass (see, *e.g.*, [1]); this is the energy needed to accelerate the whole universe outwards at its measured rate. The remaining 25 ~ 30% is Dark Matter (DM) (see, *e.g.*, [2]), and is needed to account for the proper motions of celestial bodies.

This paper investigates how averaging out inhomogeneities in matter affects the measurement of mass and energy parameters. When we calculate interactions between celestial bodies, we generally use a smooth mass density profile in galaxies or clusters; that is, we average mass distributions to simplify our calculations. However, Einstein’s theory is nonlinear, and this averaging could potentially alter what energies we expect to see. The averaged stress tensor of the real metric need not be the same as the real stress tensor of the averaged metric; this discrepancy could be pertinent to the problem of DM.

Effects of inhomogeneities and DM seem to have some characteristics in common. Smoothing of mass fluctuations should become less important at greater distances from the fluctuations; thus the effects on energy density would be greater closer to the masses, which is where halos are identified. It also make sense that the existence of halos is contingent upon some intrinsic properties of mass distributions in general, because the halos are found on such different scales (galaxies, galaxy clusters, even superclusters). This marks a difference from DE, which is seen as a constant energy density throughout the whole universe, a “cosmological constant” as Einstein originally called it.

This is not the first proposal with respect to the effects of smoothing on energy and the consequent significance for cosmology. Kolb *et al.* suggested that inhomogeneities could account for a cosmological constant and solve the problem of DE rather than DM [3]. This proposal was refuted by E. Flanagan [4] and R. Wald and A. Ishibashi [5], on the grounds that the effects of such fluctuations are far too small to compensate for the huge amounts of energy missing. In the context of DM, F. Cooperstock and S. Tieu claimed to construct a model in which a galaxy’s need for a halo was canceled by including some of its possible rotational energy [6]; M. Korzynski and others have shown this claim is erroneous [7]. Other explorations of these issues include [8].

All these calculations, however, are perturbative—based upon approximations; until now, there has not been an *exact* calculation of the impact of smoothing on energy. This paper determines the exact energy for two scenarios, one of which is a ‘smoothed’ version of the other. Both scenarios use BPS black hole solutions because it is easy to find a solution for any sum of these special black holes, as the electromagnetic repulsions and gravitational attractions between them cancel [9]. Specifically, we compare an infinite string of discrete charged black holes to a smooth line of charge, with both distributions normalized to have the same average charge per unit length. We work in five dimensions because such distributions are divergent in four dimensions.

2 The Metrics

The solution of the coupled Einstein-Maxwell field equations which describes a set of $n = 1 \dots N$ extremal black holes with charges q_n in five dimensions¹ is [10]:

$$ds^2 = -\Omega^{-2}dt^2 + \Omega d\vec{x}^2, \quad A_0 = \frac{\sqrt{3}}{2}\Omega^{-1}, \quad \Omega = 1 + \sum_{n=1}^N \frac{q_n}{|\vec{x} - \vec{x}_n|^2}; \quad (2.1)$$

as we are in five dimensions, $d\vec{x}^2 = dx^2 + dy^2 + dz^2 + dv^2$. We note that Ω is time-independent and harmonic (it satisfies $\nabla^2\Omega = 0$ outside the source), and that, in these coordinates, the horizon of each black hole is shrunk down to a point.

These are BPS solutions, with the property that gravitational attraction and electromagnetic repulsion between any set of them cancel [9]. This makes them time-independent: with no forces between them, they are in equilibrium. This is also what ensures that solutions for a set of them are easily found, namely by the sum in (2.1).

We consider two configurations: a discrete (D) and a smooth (S) solution. In (D), we have a discrete set of black holes positioned along a line at equal distances L from one another other, whereas in (S) we have smooth charged string along an axis. The continuous set can be viewed as a limiting case of the discrete set, namely the limit as the number of black holes per unit length tends to infinity and as the charge of each black hole tends towards zero, in such a way that the total charge per unit length remains the same.

2.1 The Discrete Solution (D)

For the discrete case, we choose black holes with the same charge q_D ; thus Ω_D is given by:

$$\Omega_D = 1 + \sum_{n=-\infty}^{\infty} \frac{q_D}{r^2 + (v-n)^2}, \quad (2.2)$$

where black holes are one unit length apart on the v -axis and r is the distance to the v -axis.. We can restore the separation L by dimensional analysis (keeping the total charge per unit length constant):

$$\Omega_D^L = 1 + \sum_{n=-\infty}^{\infty} \frac{Lq_D}{r^2 + (v-nL)^2} = 1 + \sum_{n=-\infty}^{\infty} \frac{q_D/L}{(r/L)^2 + (n - [v/L])^2}. \quad (2.3)$$

Thus we can start with $L = 1$ and find the dependence on L simply by making the substitution $(r, v, q_D) \rightarrow (r/L, v/L, q_D/L)$.

To evaluate the sum in (2.2), we use contour integration. We find

$$\sum_{n=-\infty}^{\infty} \frac{1}{r^2 + (v-n)^2} = \frac{i\pi \cot(\pi(v+ir))}{2r} - \frac{i\pi \cot(\pi(v-ir))}{2r}, \quad (2.4)$$

¹In four dimensions, the solution [9, 10] has the form $ds^2 = -\Omega^{-2}dt^2 + \Omega^2 d\vec{x}^2$, $A_0 = \Omega^{-1}$.

which simplifies to

$$\sum_{n=-\infty}^{\infty} \frac{1}{r^2 + (v-n)^2} = \frac{\pi}{r} \frac{\sinh(2\pi r)}{\cosh(2\pi r) - \cos(2\pi v)} . \quad (2.5)$$

Thus, using (2.2) and (2.5), the metric for the discrete solution is given by:

$$\Omega_D = 1 + \frac{\pi q_D}{r} \frac{\sinh(2\pi r)}{\cosh(2\pi r) - \cos(2\pi v)} . \quad (2.6)$$

2.2 The Smooth Solution (S)

The metric for (S) is the same as the limit of the metric of (D) as the masses and the distances between them approach zero. From (2.3), we find:

$$\Omega_S = 1 + \lim_{L \rightarrow 0} \sum_{n=-\infty}^{\infty} \frac{L q_D}{r^2 + (v-nL)^2} = 1 + \int_{-\infty}^{\infty} dx \frac{q_D}{r^2 + (v-x)^2} = 1 + \frac{\pi q_D}{r} . \quad (2.7)$$

As expected, the discrete harmonic function Ω_D (2.6) approaches the smooth function Ω_S (2.7) when the distance r to the v -axis is large.

3 Energy

Since we have normalized the two solutions so that they have the same average charge density along the v axis, and since the solutions are BPS, they must have the same total mass density as well; this does not immediately rule out a possible effect, as we discuss below.

We check this using Weinberg's definition of total energy [11]: we rewrite the metric as a flat metric plus a correction term h : $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$. We split the Einstein tensor into terms linear and non-linear in h : $G_{\mu\nu}^{(L)}(h) + G_{\mu\nu}^{(NL)}(h) = T_{\mu\nu}$. Then the definition of the total (matter plus gravitational) energy density of a system is given by [11]:

$$\text{Total Energy Density} = G_{00}^{(L)}(h) = T_{00} - G_{00}^{(NL)}(h) . \quad (3.1)$$

We integrate $G_{00}^{(L)}(h)$ over all three directions (x, y, z) orthogonal to the line of black holes. Integrating along the entire v -axis would make the integral divergent—but this is not a problem: since the smooth configuration (S) is independent of v , and the discrete configuration (D) is periodically dependent upon it, we need only integrate over one period of configuration (D) (*i.e.*, from 0 to 1 for $L = 1$). To perform the integral, we transform to 4-d cylindrical coordinates $(r, \theta, \varphi, \text{ and } v)$, and integrate over v, θ, φ , factoring in the Jacobian $r^2 \sin(\varphi)$ for the coordinate change. Then the total energy per unit length along the v -axis (describing an infinite slice of space orthogonal to the v -axis, with a width of one unit length) can be written as the following surface integral:

$$E = \int_0^1 dv \int d\theta d\varphi r^2 \sin(\varphi) n^i (\partial_i h_{jj} - \partial_j h_{ij}) , \quad (3.2)$$

where $\vec{n}$ is a unit vector normal to the v -axis. Since at large values of r the integrands are the same, this integral yields the same result for our two cases—in agreement with the requirements of the BPS condition (which ensures that interaction energies cancel) and the equality of charge per unit length (which ensures that the energies which are not due to interactions are also equal).

The equality of the total masses does not in itself preclude the possibility of an effect on DM: the energy which enters Einstein's equations, and so is relevant to DM, is the covariant matter energy T_{00} . This quantity is not necessarily the same for our two different distributions of black holes; comparing them shows us how matter energy is affected by smoothing.

3.1 Stress Tensor and Integration

The matter energy of a system is given by the integral of T_{00} over all space. In a general spacetime with a time-like Killing vector ξ^ν the following total derivative vanishes:

$$0 = \int d^5x \sqrt{-g} D_\rho (g^{\rho\mu} T_{\mu\nu} \xi^\nu) = \int d^5x \partial_\rho (\sqrt{-g} g^{\rho\mu} T_{\mu\nu} \xi^\nu) ; \quad (3.3)$$

consequently, the contribution on a spacelike slice defines a conserved energy. In our case, since both the (D) and (S) metrics are static and diagonal, the total matter energy reduces to:

$$E = - \int d^4x \sqrt{-g} g^{00} T_{00} . \quad (3.4)$$

The stress-tensor is usually defined with a factor $\frac{1}{4\pi}$, but as we only study ratios of energies, we are not interested in the overall normalization.

We calculate T_{00} using

$$T_{\mu\nu} = F_{\mu\alpha} F_{\nu\beta} g^{\alpha\beta} - \frac{1}{4} g_{\mu\nu} (F_{\alpha\beta} F_{\gamma\delta} g^{\alpha\gamma} g^{\beta\delta}) , \quad F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu , \quad (3.5)$$

where A is the gauge potential (2.1). Once again, because the metrics are static and diagonal, and furthermore, because the gauge potential A is purely timelike and static, this simplifies, and we find

$$T_{00} = F_{0j} F_{0j} g^{jj} - \frac{1}{4} g_{00} (2 F_{0j} F_{0j} g^{00} g^{jj}) . \quad (3.6)$$

Recalling $A_0 = \frac{\sqrt{3}}{2} \Omega^{-1}$, $g_{00} = -\Omega^{-2}$, $g_{jj} = \Omega$, and the determinant $g = -\Omega^2$ from equation (2.1), we find:

$$E = - \int d^4x \sqrt{-g} g^{00} T_{00} = 6 \int d^4x \left(\frac{\partial_j \Omega}{\Omega} \right)^2 . \quad (3.7)$$

Here $(\partial_j \Omega)^2 = (\partial_x \Omega)^2 + (\partial_y \Omega)^2 + (\partial_z \Omega)^2 + (\partial_v \Omega)^2$.

3.2 Exact Energies

As discussed above equation (3.2), we integrate stress-energy over all three directions (x, y, z) orthogonal to the line of black holes and integrate over one period of configuration (D) over the v -axis. The smooth configuration (S) is the simpler of the two; we calculate it first. As discussed in section 2.1, we use $L = 1$, and reintroduce L later by rescaling. We ignore integration over θ and φ in both configurations because it only produces equal overall constants; we also drop the $\sin(\varphi)$ factor from the Jacobian. Defining $k = \pi q$ and $\Omega_S = 1 + \frac{k}{r}$, we find $(\partial_j \Omega_S)^2 = k^2/r^4$, and then (3.7) gives

$$E_S = \int_0^\infty dr \frac{k^2}{(r+k)^2} = k. \quad (3.8)$$

For discrete configuration (D) , the calculation is more complicated. We define $s = \sinh(2\pi r)$, $c = \cosh(2\pi r)$, $t = \sin(2\pi v)$, and $d = \cos(2\pi v)$ and compute:

$$(\partial_j \Omega_D)^2 = \frac{k^2 [s(d-c) + 2\pi r(1-cd)]^2 + (2\pi rst)^2}{r^4 (c-d)^4}. \quad (3.9)$$

Then, with $\Omega_D = 1 + \frac{k}{r} \frac{s}{c-d}$, and the measure factor r^2 , the energy for (D) is given by:

$$E_D = \int_0^1 dv \int_0^\infty dr k^2 \frac{[s(d-c) + 2\pi r(1-cd)]^2 + (2\pi rst)^2}{(c-d)^2 (r(c-d) + ks)^2}. \quad (3.10)$$

This is not easily evaluated as an indefinite integral, but we can compare E_S and E_D numerically. To understand the results, let us clarify what we should test. For $L = 1$ and various k (charge), we want to find the differences in energy between configurations (D) and (S) . However, a smaller k will certainly yield a smaller energy and smaller energy discrepancy, so to compare various k 's we divide by E_S because $k \propto E_S$.

Thus we study the ratio $\left(\int_0^\infty \int_0^1 (E_D - E_S) dv dr \right) / \left(\int_0^\infty \int_0^1 E_S dv dr \right)$ as a function of k . Different k represent different densities, as we have scaled the separation L to $L = 1$. The results, computed with Maple, are consistent with zero; the actual numerical integrations are somewhat delicate because of divergences in the integrand (E_D diverges when $r = 0$ and $v = 0, 1$).

4 Physical Application

Though we did not find an effect, we discuss how any effect that we might have found could have been interpreted physically. The first step would have been to choose a scale, because DM is found on several different levels: galaxies, galaxy clusters, superclusters. Then we would determine which objects are smoothed for the situation chosen. For example, to see how smoothing might create a halo around a galaxy, we would smooth the stars inside the galaxies, as is the common practice when calculating intergalactic motions.

Our numerical results were calculated for $L = 1$. We can use these results, however, if we rescale the situation by changing the mass so that the density of the galaxy is correct. The value of

k in equations (3.8) to (3.10) would then be modified to provide the correct mass matching up with the real-life situation. The k we used was πq , and $q = Gm$ for extremal black holes. The ratio to demand for similarity is $k/1 = \pi Gm/L$.

So, we need the mass (m) and average distance between masses (L) as input, and then we can see what the energy difference is for $k = \pi Gm/L$. Let us apply these ideas to the situation of galaxy halos just discussed, testing how the smoothing of stars within the galaxies could affect the energy. We use planck-units, so that $G = 1$. The average distance between adjacent stars is about 10 light-years, 10^{52} Planck lengths. The average mass of a star is a bit less than 10^{39} Planck masses. Our k is then around 10^{-13} .

Next let us apply the smoothing to galaxies within galaxy clusters. The average galaxy mass is 10^{45} Planck-masses. The average separation between galaxies is 10^7 light-years, 10^{58} Planck-lengths. So k for this situation is again 10^{-13} .

A third scale at which to look is that of superclusters. Superclusters consist of about a dozen clusters, each with a mass of about 10^{55} Planck-masses. Usually they are arranged in strings with lengths of 10-100 Mpc, 10^{58} or 10^{59} Planck-lengths, so the distance between each is about 10^{57} or 10^{58} . Our k is then 10^{-2} or 10^{-3} .

5 Conclusion

The calculation performed in this paper is by no means flawless. Five-dimensional effects are expected to show similar trends as their four-dimensional counterparts, but when one is discussing the extent of an effect, 5-d is less applicable. Also, these calculations were done with BPS solutions. They are easy to work with but certainly do not represent standard matter. They are maximally charged, for one thing, and they describe point-like masses, a physical impossibility.

These are the first exact calculations of the effects of inhomogeneities; as such, they complement and confirm perturbative arguments. There is room for development—our calculation was based on a particular model, and calculations in other models could help elucidate the exact effects of smoothing.

Acknowledgements

We are happy to thank Peter van Nieuwenhuizen for extensive discussions, encouragement, and comments. We also thank Nick Jones for comments. MR thanks the Institute for Theoretical Physics at the University of Amsterdam for hospitality. MR is supported in part by NSF grant no. PHY-0354776, by the University of Amsterdam, and by Stichting FOM.

References

- [1] S. Perlmutter *et al.* [Supernova Cosmology Project Collaboration], *Measurements of Omega and Lambda from 42 High-Redshift Supernovae*, *Astrophys. J.* **517**, 565 (1999) [arXiv:astro-ph/9812133].
- U. Seljak *et al.*, *Cosmological parameter analysis including SDSS Ly-alpha forest and galaxy*

- bias: Constraints on the primordial spectrum of fluctuations, neutrino mass, and dark energy*, Phys. Rev. D **71**, 103515 (2005) [arXiv:astro-ph/0407372].
- [2] G. Bertone, D. Hooper and J. Silk, *Particle dark matter: Evidence, candidates and constraints*, Phys. Rept. **405**, 279 (2005) [arXiv:hep-ph/0404175].
- [3] E. Kolb and S. Matarrese and A. Riotto. *On cosmic acceleration without dark energy*, arXiv:astro-ph/0506534.
- [4] E. E. Flanagan, *Can superhorizon perturbations drive the acceleration of the universe?*, Phys. Rev. D **71**, 103521 (2005) [arXiv:hep-th/0503202].
- [5] R. Wald and A. Ishibashi. *Can the Acceleration of Our Universe Be Explained by the Effects of Inhomogeneities?*, arXiv:gr-qc/0509108.
- [6] F.I. Cooperstock and S. Tieu. *General Relativity Resolves Galactic Rotation Without Exotic Dark Matter*. arXiv:astro-ph/0507619.
- [7] M. Korynski. *Single disk of matter in the Cooperstock-Tieu galaxy model* arXiv:astro-ph/0508377.
D. Vogt and P.S. Letelier. *Presence of exotic matter in the Cooperstock and Tieu galaxy model* arXiv:astro-ph/0510750.
D. Garfinkle, *The need for dark matter in galaxies*, arXiv:gr-qc/0511082.
- [8] T. Buchert, *On globally static and stationary cosmologies with or without a cosmological constant and the dark energy problem*, arXiv:gr-qc/0509124.
G. F. R. Ellis and T. Buchert, *The universe seen at different scales*, Phys. Lett. A **347**, 38 (2005) [arXiv:gr-qc/0506106].
T. Buchert and M. Carfora, *The Cosmic Quartet - Cosmological Parameters of a Smoothed Inhomogeneous Spacetime*, arXiv:astro-ph/0312621.
T. Buchert and M. Carfora, *Cosmological parameters are dressed*, Phys. Rev. Lett. **90**, 031101 (2003) [arXiv:gr-qc/0210045].
T. Buchert, M. Kerscher and C. Sicka, *Backreaction of inhomogeneities on the expansion: the evolution of cosmological parameters*, Phys. Rev. D **62**, 043525 (2000) [arXiv:astro-ph/9912347].
- [9] S.D. Majumdar, Phys. Rev. **72**, 930 (1947).
A. Papapetrou, Proc. R. Irish Acad. **A51**, 191 (1947).
- [10] G. W. Gibbons, D. Kastor, L. A. J. London, P. K. Townsend and J. H. Traschen, *Supersymmetric selfgravitating solitons*, Nucl. Phys. B **416**, 850 (1994) [arXiv:hep-th/9310118].
- [11] S Weinberg. *Gravitation and Cosmology : Principles and Applications of the General Theory of Relativity*, Chapter 7, Section 6. Wiley, New York, 1972.

- [12] G. Tormen and F.R. Bouchet and S.D.M. White. *The Structure and Dynamical Evolution of Dark Matter Halos*. arXiv:astro-ph/9603132